This Page Is Inserted by IFW Operations and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents will not correct images, please do not report the images to the Image Problem Mailbox.

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 5: (11) International Publication Number: WO 92/04009 A1 A61K 9/127 (43) International Publication Date: 19 March 1992 (19.03.92)

(21) International Application Number: PCT/GB91/01547 (74) Agent: GILL JENNINGS & EVERY; 53/64 Chancery Lane, London WC2A 1HN (GB). (22) International Filing Date: 10 September 1991 (10.09.91)

(71) Applicant (for all designated States except US): SCHOOL OF PHARMACY, UNIVERSITY OF LONDON [GB/ GB]; 29/39 Brunswick Square, London WC1N 1AX

10 September 1990 (10.09.90) GB

10 September 1990 (10.09.90) GB

(GB).

(72) Inventor: and (75) Inventor/Applicant (for US only): GREGORIADIS, Gregory [GB/GB]; "Crantock", Kewferry Hill, Rickmansworth Road, Northwood, Middlesex HA6 2RQ (GB).

(81) Designated States: AT (European patent), BE (European patent), CH (European patent), DE (European patent), DK (European patent), ES (European patent), FR (European patent), GB (European patent), GR (European patent), IT (European patent), JP, LU (European patent), NL (European patent), SE (European patent), US.

Published

With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(54) Title: LIPOSOMES

(30) Priority data:

9019726.0

9019729.4

(57) Abstract

Novel liposomes have a mean diameter not exceeding 200 nm and containing at least 33 mg of one or more entrapped substances per g liposome-forming material. They may also comprise, co-entrapped or covalently linked in the same liposome, a synthetic peptide and an adjuvant which potentiates the immunogenicity of the peptide. The liposomes are useful in vaccines.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphiets publishing international applications under the PCT.

AT	Austria	· E5	Spain	MC	Moderascar
AÜ	- Australia	PI	Finland	ML.	Mili
28	Barhados .	FR	France	MN	Mongolia
BE	Belgium	GA	Gabon	MR	Mauritania
BP	Burkina Faso	GB	United Kingdom	MW	Malawi Malawi
BC	Bulgaria	GN	Guinca	NL NL	Netheriands
BJ	Benin	GR	Gruce		
BR	Brazil	HU		NO	Norway
CA	Canada	IT -	Hungary	PL	Poland
CF	Central African Republic		luly	RO	Romania
œ	Congo	JP	Japan	SD	Sudan
CII	Switzerland	KP	Democratic People's Republic	SE	Sweden
_			of Korea	SN	Sencyal
Cl	Côte d'Ivoire	KR	Republic of Korea	SU+	Soviet Union
СМ	Cameroon	LI	Liechtenstein	TD	Chad
CS	Czechoslovakia	LK	Sci Lanka	TC	Togo
DE•	Germany	LU	Luxembourg	US .	United States of America
DK	Denmark	MC	Manua		

⁺ Any designation of "SU" has effect in the Russian Federation. It is not yet known whether any such designation has effect in other States of the former Soviet Union.

35

1

LIPOSOMES

Field of the Invention

The present invention relates to liposome preparations which combine small size with very high retention of added substances. The present invention also relates to liposomes containing synthetic peptides, e.g. for use in vaccines.

Background of the Invention

High efficiency entrapment (retention) of substances using minimal amounts of lipid is a prerequisite for the 10 effective use of liposomes in a variety of applications. For example, a high entrapped drug to lipid mass ratio is necessary for an acceptable cost of formulation and an acceptable low risk of lipid-induced toxicity 15 pharmaceutical use. Unfortunately, current liposome preparations and methodologies giving high efficiency entrapment, as described by Szoka et al, Proc. Nat. Acad. Sci. USA 75 (1978) 4194-4198, and by Kirby and Gregoriadis, Biotechnology 11 (1984) 979-984, result in large liposomes, having mean diameters approaching 1 μm in size. This large 20 size severely limits, for example, pharmaceutical use: such large liposomes, on intravenous injection, exhibit short half-lives in blood circulation; see Senior et al, Biochim. Biophys. Acta <u>839</u> (1985) 1-8. They are therefore unsuitable for maintaining increased concentrations of 25 drugs and other substances within the vascular system. In addition, such large liposomes are unsuitable for use as delivery systems for targeting of drugs or other substances to specific sites, for example to diseased cells in the 30 body.

Smaller liposomes, 200 nm or less in diameter, are considerably more stable, for example in the bloodstream, than the larger liposomes described above. Other features of smaller liposomes for pharmaceutical applications are that, unlike larger liposomes, they are able to reach the lymph nodes efficiently and to deliver vaccines and other agents to these sites: see Senior et al, loc. cit.; Turner

35

et al, Biochim. Biophys. Acta 760 (1983) 119-125; Allen et al, FEBS Letters 223 (1987) 42-46; Chow et al, Pharmacol. Exper. Ther. 248 (1989) 506-513; and Spanzer et al, Biochim. Biophys. Acta 836 (1986) 224-230. However, current techniques for preparing small liposomes, as described by Mayhew et al, Biochim. Biophys. Acta 775 (1984) 169-174, suffer from either a low efficiency of entrapment of substances or from a large lipid requirement (180 μ mol per ml or greater) for efficient entrapment.

Most current vaccine preparations for infectious 10 disease comprise either attenuated or killed infectious Such preparations suffer from problems such as safety, for example reversion of attenuated to virulent and lack of effectiveness, for example an strains. 15 incidence non-responsive recipients. of widespread interest in the possibility of synthetic peptide vaccines, especially against viral, bacterial and parisitic diseases (see Arnon, Synthetic Vaccines, 1987 (vols. 1 and 2), CRC Press Inc.) and, more recently, in cancer. The principle of a synthetic peptide vaccine is that the 20 protective epitopes on, for example, infectious agents are copied as short synthetic peptides free of any infectious agent. In order to be effective, a synthetic peptide must possess a high level of immunogenicity and antibodies which cross-react extensively with the target 25 organism or cell; see Van Regenmortel, Immunology Today 10 (1989) 266-272. In practice, peptides alone tend to possess weak immunogenicity, and anti-peptide antibodies often do not recognise the native protein antigen; see Jemmerson et al, Molecular Immunology 26 (1989) 301-307. 30

A common method for enhancing the immunogenicity of synthetic peptide epitopes is to combine the peptides with an immunological adjuvant. Adjuvants are agents which provoke potent immune responses to antigens, and include aluminium hydroxide (alum), saponins, pluronic polymers with mineral oil, killed mycobacterium in mineral oil (Freund's complete adjuvant), bacterial products such as

10

15

25

muramyl dipeptide, and immunogenic carrier proteins such as tetanus toxoid. Adjuvants, depending on their nature, induce humoural (HI) and/or cell-mediated immunity (CMI). However, most of such adjuvants are either unsuitable for combination with peptides, have toxic side-effect problems, or fail to induce both HI and CMI; see Gregoriadis, Immunology Today 11 (1990) 89-97. A further problem is interference of peptide immunogenicity by immune response to a carrier protein; see Etlinger et al, Science 249 (1990) 423-425.

As described elsewhere (see Gregoriadis, loc. cit.), liposomes have the potential for a safe and effective adjuvant for peptide vaccines, and the immunopotentiation of a number of synthetic peptides has been shown. However, such immunopotentiation by liposomes is not a general phenomenon. For example, liposomal immunopotentiation has not been observed with a pre-S peptide (amino acids 15-48) from Hepatitis B surface antigen (HBsAg) or with a 3VP2 peptide from poliovirus.

Summary of the Invention 20

Novel, small liposomes have a mean diameter less than 200 nm and high entrapment efficiency, and require low (<150 μ mols per ml) concentrations of starting lipid. They are suitable for use in a variety of applications, including, for example, cosmetics for the sub-epidermal delivery of substances by liposomes, medical diagnostics for the stable entrapment of easily measurable substances, and pharmaceuticals (as described above).

Further, novel synthetic peptide vaccine preparations comprise liposomes containing entrapped or covalently-30 linked peptides together with molecules which augment the immune response to the peptides.

Description of the Invention

Examples of substances that may be entrapped in liposomes of the invention are pharmaceuticals, vaccines, 35 materials, enzymes, hormones, vitamins, carbohydrates, proteins/peptides, lipids, organic

4

molecules, and inorganic molecules or atoms. More specific examples are anti-tumour and anti-microbial agents, enzymes, hormones, vitamins, metal chelators and genetic material, preferably carbohydrates or proteins/peptides. The liposomes may be incorporated into vaccines.

The liposomes may be produced from one or more of phosphatidylcholine, cholesterol, phosphatidylglycerol, phosphatidylethanolamine, phosphatidylserine, phosphatidic sphingomyelin or derivatives of these lipids, preferably phosphatidylcholine or cholesterol. The process 10 for their preparation may be from a starting lipid concentration of less than 150 μ mols per ml, to provide an efficiency of entrapment of >10% of a starting substance concentration. The method involves a combination of the dehydration-rehydration vesicle (DRV) process (see Kirby 15 and Gregoriadis, loc. cit.) and the microfluidation process described by Talsma et al, Drug Development Ind. Pharm. 15 (1989) 197-207. Briefly, the method involves mixing lipids with substances to be entrapped and subjecting the mixture to controlled dehydration and rehydration to form large 20 multilamellar vesicles with high entrapment efficiency. These large vesicles are then subject to microfluidation to produce small liposomes with 10-100% retention of the entrapped substances.

25 As will be apparent from the evidence given below, microfluidisation of solute-containing DRV's in PBS and the presence of unentrapped solute produces vesicles less than 200 nm in size, and which are suitable for the given purposes. Such vesicles retain, under the present conditions, 35-78% of the originally entrapped solute. 30 Compared to procedures (Mayhew et al, loc. cit.) which employ much larger amounts of lipid to achieve efficient entrapment, the present approach provides preparations with augmented solute to lipid mass ratios and should also be more economical. The narrow distribution of final vesicle 35 sizer must reflect the nature of the production process. Whereas sonication disrupts vesicles in a random mode,

20

25

producing a wide range of diam ers, microfluidisation distorts large and flexible vesicles by extrusion through a capillary, thereby increasing the number of vesicles of similar size which "bud off" from simple parent vesicles.

While being applicable across the broad range of the invention, the minimum amount of retained solute is 35%, i.e. 2.5 mg with respect to a starting amount of 7.2 mg of the relevant substance. This is for up to 66 µmoles, i.e. up to 50 mg, phosphatidylcholine (approximate molecular weight 760) up to 25 mg cholesterol (molecular weight 387). Therefore, by means of the present invention, it is possible to produce liposomes having mean diameters of less than 200 nm which contain at least 50 mg of the entrapped substance per g phospholipid or at least 33 mg of the substance per g of liposome-forming materials. In Example 1, the liposomes retain approximately 35-78% of the originally entrapped test substances maltose or tetanus toxoid (herein referred to as solutes) following microfluidisation. This corresponds to an overall solute entrapment yield of >10%.

For example when including one or more adjuvants, liposome preparations are made by standard methods for producing liposomes, for example as described by Gregoriadis, ed., Liposome Technology (1984) vols. 1 to 3, CRC Press Inc. In such methods, liposome components, peptides and adjuvant molecules are either mixed or covalently linked prior to formation of liposomes.

The following Examples illustrate the invention, with reference to literature articles given above and also to Deamer and Uster, Liposome Preparation: Methods and mechanisms in: ed. Ostro, Liposomes, M-tcel Dekker, Inc., New York, 1983, pp 27-51; Gregoriadi. et al, Vaccine 5 (1987) 145-151; and Davis and Gregoriadis, Immunology 61 (1987) 229-234.

The source and grades of egg phosphatidylcholine (PC) and cholesterol have been reported elsewhere (Kirby and Gregoriadis, 1984). Maltose was from Sigma Chemical

Company (London) and immunopurified tetanus toxoid from Wellcome Biotechnology (Beckenham). D-[U- 14 C]-maltose (20 GBq per μ mol) was purchased from Amersham International (Amersham, U.K.) and toxoid-labelled (specific activity 5.43 mBq) with 125 I as described elsewhere (Gregoriadis et al, 1987).

The following abbreviations are used:

60 psi = 414 kPa

PBS - Phosphate-Buffered Saline

10 PC - Phosphatidylcholine

DRV - Dehydration-Rehydration Vesicles

SUV - Small Unilamellar Vesicles

dpm - disintegration per minute

Example 1

15 <u>Preparation of liposomes</u>

Solute-containing DRV liposomes were prepared by the procedure discussed elsewhere (Kirby and Gregoriadis, 1984; Gregoriadis et al, 1987). Briefly, small

unilamellar vesicles (SUV) made in distilled water from equimolar PC (16.5, 33, 65 or 132 µpol) and cholesterol were mixed with an equal volume (2.0 ml) of maltose (7.2 mg) or tetanus toxoid (7.2 mg) dissolved in PBS (0.44mM sodium phosphate, 2.7mM potassium chloride and 0.14 M sodium chloride, pH 7.4) and supplemented with traces (4.9 \times 10 - 8.7 \times 10 dpm) of the respective rediclabelled solutes. Following dehydration and controlled (Kirby and Gregoriadis, 1984) rehydration to form solute-containing DRV, the suspensions (2.0 ml) were each divided into two equal portions. One portion was centrifuged at 20,000g for 30 min, the pellet washed three times with PBS and suspended in 20 ml of the same buffer (washed DRV) for microfluidization. Entrapment of solutes was estimated by the assay (Kirby and Gregoriadis, 1984; Gregoriadis et al, 1987) of ¹⁶C or ¹⁸I radioactivity in the suspension, and results expressed as percentage of the solute originally present. The other portion (containing a mixture of entrapped and unentrapped solute) was also diluted in 20 ml PBS (unwashed DRV) and then microfluidized. In some experiments, PBS was replaced by distilled water in all steps involved in the entrapment of solutes in DRV and microfluidization of washed or unwashed preparations.

Microfluidization of liposomes

Washed and unwashed DRV (20 ml each) were passed for 1.8, 3.5, 5.2, 7.1 and 10.6 full cycles through a Microfluidizer 110[™] kindly provided by Microfluidics Corp., Newton, MA, USA. The pressure gauge was set at 60 psi throughout the procedure to give a flow rate of 35 ml per minute. At the end of each of the 20 ml cycle intervals, samples containing maltose were dialysed exhaustively

against distilled water or PBS, and the toxoid containing vesicles or centrifuged for 30 min at 20,000g (1.8 and 3.5 cycles) or 35,000g (5.2, 7.1 and 10.6 cycles), the pellets washed twice in PBS or water and resuspended in 1.0 ml of the respective solvents. In the case of the toxoid, the supernatants obtained on centrifugation were passed (Senior et al. 1985) through Sepharose 4B columns; very little (less than 3%) of the solute in the centrifuged samples eluted in the liposome form. The extent to which washed and unwashed DRV's retained their solute content after microfluidization was estimated as follows: for washed DRV'S, radioactivity retained after dialysis or recovered in pellets after centrifugation was expressed as a percentage of the total amount in the sample before dialysis or centrifugation. Solute retention values thus obtained correspond to the percentage of the original entrapment values in the preparations before microfluidization. For unwashed DRV, the percentage of radioactivity retained after dialysis or centrifugation was expressed as % of the original entrapment values.

Measurement of vesicle size

Particle size distributions were measured by photon correlation spectroscopy of samples diluted in either water or PBS, using a Malvern Model 4700 apparatus (Malvern Instruments Ltd., Malvern, UK) equipped with a 25mW helium/neon laser. Mean diameters and size distributions are obtained: the z-average mean diameter, polydispersity factor and cumulative percentage mass and number undersize data were recorded as a function of the number of microfluidization cycles. The performance of the instrument was checked with monodisperse

polystyrene latex suspensions (Polysciences, UK) and mixtures of such latex samples to verify the ability of the system to accurately measure polydisperse or bidisperse systems.

Results and Discussion

The effect of DRV microfluidization on solute retention

As expected from previous work (Kirby and Gregoriadis, 1984; Gregoriadis et al 1987), entrapment of solutes in dehydration-rehydration liposomes was efficient, ranging from 19.2 to 66.1% of the starting material and dependent on the amount of phospholipid used (see legends to Figs 1-3). There was no significant difference in entrapment values in preparations made in distilled water or PBS (not shown). Retention values (quoted as a percentage of original entrapment values) by solute-containing DRV's microfluidized for up to 5.2 cycles in the presence of PBS are shown in Figs 1 and 2. It is apparent that retention values by both washed and unwashed DRV decrease or remain the same as the number of cycles increases (see also Fig. 3). Thus, values of 80-100% (maltose) after 1.8 cycles become 40-65% after 5.2 cycles (Fig. 1), whereas values for the toxoid (unwashed DRV) remain virtually unchanged (at around 75%) (Fig. 2). It is conceivable that during microfluidization, a process entailing breakage of the integrity/vesicle membrane and consequent leakage of entrapped solute occurs, large molecules such as proteins leak at slower rates than smaller molecules. Furthermore, no apparent difference in solute retention was observed for preparations containing varying amounts of PC (8.5-66 µmoles) (Figs. 1 and 2). Figs. 1 and 2 also indicate that, generally, a greater proportion of solute (maltose or toxoid) is retained by unwashed vesicles (ie. preparations microfluidized in the presence of unentrapped solute) than by washed DRV. The presence of unentrapped solute during microfluidization diminishes solute leakage perhaps by reducing the osmotic rupture of the vesicles and by reducing initial concentration gradients across the membrane.

The mean diameter of microfluidized DRVs measured by dynamic light scattering after a given cycle interval depended on whether the DRVs were suspended in distilled water or PBS. We therefore attempted to determine the effect of the two media on solute retention during microfluidization. Results in Fig. 3 indicate that much more maltose is retained by DRVs in the presence of PBS and similar results (not shown) were obtained with toxoid-containing DRV's under identical conditions. Such an effect of PBS presence in the milieu could be attributed to its reduction of osmotic shock expected (Kirby and Gregoriadis, 1984) to occur on dilution of solute-containing DRV in hypotonic media.

The effect of microfluidization on vesicle size

The mean diameters of washed and unwashed maltose-containing DRV microfluidized in water or PBS (from experiment described in Fig. 3), measured by photon correlation spectroscopy, are presented in Table 1. They show that, in agreement with findings by others (Talsma et al, 1989), a smaller vesicle size was achieved for preparations processed in water than in PBS. Such differences in sizes, appearing after the 1.8 cycle interval, could be attributed (Deamer and Uster, 1983) to vesicle aggregation induced by the presence of salts in the PBS

medium, but also might be due to the increased lipophilicity of the lipids in the presence of electrolyte. Reduction in the size of all preparations tested was considerable after the initial I.8 cycles interval. Subsequently, sizes levelled off to reach minimal values of 100-160 nm by 10.6 cycles depending on the medium (water or PBS) (it is of interest that a pattern similar to that of sizes in Table 1 was observed for solute retention with increasing cycle intervals (Fig.3), with values also levelling off after 3.5 cycles). Mean number diameter and mass diameter distribution data (Fig. 4 and Fig. 5) from the experiments recorded in Table I revealed a bimodal distribution of sizes in which there is an increasing number of small vesicles as the number of microfluidization cycles increases beyond the first few cycles; the mean number diameter of the sample, prepared in water (50mg maltose/water) is of the order of 300mm after 1.8 cycles, with a few vesicles with diameters up to lum in diameter. The larger diameter vesicles disappear rapidly on increasing cycling, so that after 10.6 cycles in water, a single narrow diameter (around 100 nm) distribution is obtained whereas in PBS the bimodal pattern is retained (Fig 4). However, the vesicle size distribution does not exhibit the gradual downward drift that occurs on, say, sonication but a transfer of sizes from one size band to another. In PBS, some vesicles undoubtedly flocculate so that there remains a percentage of vesicles with a high apparent diameter. Fig. 6 illustrates the distribution of diameters by mass. Here is clear evidence of the marked narrowing of the polydispersity of the vesicles. Coupled with the data from Fig. 4, this figure clearly shows the narrow distribution of sizes which is achieved after 10 cycles. Lidgate and coworkers' (1989) results show that in preparing parenteral emulsions of squalene by microfluidization, the particle size after 9-10 cycles is in the range of

0.07 to 0.2 μm which represent a much narrower distribution than other technologies.

Table 1 shows the average mean size (nm) microfluidised DRV, i.e. maltose-containing washed or unwashed DRV (33 μ moles) PC were microfluidised in the presence of water or PBS for up to 10.6 cycles and samples measured for vesicle size (diameter in nm) by dynamic light scattering (Photon correlation spectroscopy). Polydispersity indexes ranging from 0.503 to 0.653 (water) and 0.517 to 0.653 (PBS) were similar to those obtained with some of the lipid compositions of liposomes employed by Talsma et al (1987).

Table 1. z - Average mean size (nm) of microfluidised DRV

			Cycles		
DRV	1.8	3.5	5.2	7.1	10.
Washed Water PBS	463.5 447.4	149.9 198.6	115.0	121.9 159.5	114 155
Unwashed	4.7.0			233,3	133
Water PBS	473.9 456.3	132.9 186.2	116.9 186.8	116.6 169.8	101 159

10

10

15

30

Description of the Drawings

Figure 1 shows entrapment values (on a scale of 0-100% of originally entrapped starting material) for 4C-labelled maltose by DRVs following microfluidisation, with respect to cycles on the abscissa. "C-maltose containing DRVs composed equimolar phosphatidylcholine of cholesterol were microfluidised in the washed (filled bars) or unwashed (shaded bars) forms in the presence of PBS for up to 5.2 cycles. Amounts of liposomal phospholipid passed through the microfluidiser were 8.25 (A), 16.5 (B), 33 (C) and 66 (D) μ mols. Values denote % retention of the originally entrapped 14C-maltose. Original entrapment values were 27.1 (A), 33.6 (B), 66.1 (C) and 65.7 (D).

Figure 2 shows entrapment values for 125 I-labelled tetanus toxoid by DRVs following microfluidisation. Details are as in Figure 1 except that amounts of liposomal phospholipid passed through the microfluidiser were 8.25 (A) and 33 (B) μ mols and original entrapment values were 19.2 (A) and 56 (B).

Figure 3 shows the effect of the liquid medium on ¹⁴Cmaltose retention by DRVs during microfluidisation.

Drtails are as in Figure 1 except that DRVs were
m_crofluidised 1. the presence of distilled water (A) or
PBS (B) for up to 10.6 cycles, the amount of liposomal
phospholipid passed through the microfluidiser was 33 μmols
and original entrapment values were 54.3 (A) and 55.2 (B).

Figure 4 shows the size distribution (as a percentage number) of vesicles containing maltose prepared in the presence of water as a function of the number of cycles ($\Delta = 10.6$ cycles; = 3.5 cycles; 0 = 1.8 cycles) through the microfluidiser, showing progression to a mean distribution with a mean diameter (on the abscissa) of approximately 100 nm after 10.6 cycles.

Figure 5 is on the same basis as Fig. 4. It shows the size distribution of vesicles containing maltose prepared in the presence of PBS (50 mg maltose/PBS) as a function of the number of cycles through the microfluidiser, showing

that, after 10.6 cycles, the number of vesicles with diameters greater than 300 nm is decreased but a biphasic distribution is still maintained.

Figure 6 shows the mass distribution of vesicles containing maltose (50 mg maltose/water) as a function of cycles through the microfluidiser as in Figure 4, showing the narrow distribution size after 10.6 cycles.

Example 2

This Example shows that interleukin-2 (IL-2) acts as an effective adjuvant for the poliovirus 3-VP2 peptide when 10 entrapped with the peptide in the same liposome. mice were injected intramuscularly in groups of five on days 0 and 28 with either 5 μ g free 3-VP2 peptide, or with liposomes containing 5 μg peptide, or with liposomes containing 5 μ g peptide and 7000 Cetus units of IL-2 15 (recombinant mutein IL-2, obtained from Cetus). Liposomes were formed from equimolar distearoyl phosphatidylcholine and cholesterol using the dehydration-rehydration procedure (see Kirby and Gregoriadis, 1984). Ten days following the second injection, serum IgG antibodies specific for 3-VP2 20 peptide were estimated by ELISA assay (see Davis and Gregoriadis, 1987) whereby microtitre wells were coated with 3-VP2 peptide, diluted sera samples (1:1280 for IgG1, 1:160 for other subclasses) were added, and bound IgG's were measured through binding of IgG subclass specific 25 antibodies. The ELISA signal measured as optical density at 492 nm was produced using horseradish peroxidaselabelled second-antibody and 3,3',5,5'-tetramethylbenzidine (Miles) as substrate. Table 2 gives the results of the IgG subclass ELISA (median values underlined). Comparison by 30 Kruskall-Wallis statistical analysis of values for groups of animals receiving liposomal 3-VP2 and liposomal 3-VP2/IL-2 revealed a significant elevation of IgG levels due to the effect of IL-2 for all IgG subclasses except IgG2b. 35 Example 3

SJL H-2' mice were injected intramuscularly in groups of five on days 0 and 28 with either 20 μg free pre-S

15

20

peptide or with liposomes containing 20 μ g pre-S peptide or with liposomes containing 20 μg S peptide and 20 μg pre-S Liposomes were formed from equimolar egg phosphatidylcholine and cholesterol using the dehydrationrehydration procedure (see Kirby and Gregoriadis, 1984). 5 Ten days following the second injection, serum IgG antibodies specific for pre-S peptide were estimated by ELISA assay (see Davis and Gregoriadis, 1987), whereby microtitre wells were coated with pre-S peptide, diluted samples (1:640) were added, and bound IgG; was measured through binding of IgG, specific antibodies. signal measured as optical density at 492 nm was produced using horseradish peroxidase-labelled second-antibody and 3,3',5,5' tetramethylbenzidine (Miles) as substrate. Table 3 gives the results of the IgG, subclass by ELISA (median values underlined).

Statistical analysing by Kruskall-Wallis revealed highly significant increase in IgGI response to pre-S peptide for liposomes with co-entrapped pre-S and S peptides.

Peptide	Secondary immune response				
formulation	IgG _l	IgG _{2s}	IgC _{Zb}	IgG3	
Free 3VP2	0.086	0.074	0.075	0.125	
	0.092	0.085	0.079	0.128	
	0.095	0.086	0.080	0.135	
•	0.096	0.113	0.080	0.141	
	0.099	0.180	0.084	0.149	
Liposomal 3VP2	0.110	0.100	0.079	0.151	
	0.112	0.107	0.085	0.161	
	0.123	0.124	0.093	0.185	
	0.130	0.143	0.096	0.206	
	0.181	0.144	0.096	0.277	
Liposomal 3VP2 +					
IL-2	0.101	0.101	0.070	0.153	
	0.285	0.143	0.075	0.269	
	0.341	0.334	0.076	0.323	
	0.448	0.492	0.104	0.359	
	1.021	0.935	0.142	0.499	
Kruskall-Wallis	H=10.5	Н=5.95	H=3.64	H=10.5	
(one-tay)	p<0.009	0.01 <p<0.049< td=""><td>p>0.102</td><td>p<0.009</td></p<0.049<>	p>0.102	p<0.009	

Peptide formulation

17

Secondary immune response to pre-S peptide

IgG₁

Free pre-S peptide	0.080	
	0.090	
	C . 095	
. •	(100	
	0.100	
Liposomal pre-5 peptide	0.075	
	0.080	
	0.085	
	0.090	· •
	0.110	
Liposomal pre-S and	0.610	
S peptides	0.680	
• •	0.780	
•	0.850	
•	0.900	

TABLE 3.

CLAIMS

- 1. Liposomes having a mean diameter not exceeding 200 nm and containing at least 33 mg of one or more entrapped substances per g liposome-forming material.
- 5 2. Liposomes according to claim 1, wherein the one or more substances is covalently linked to a lipid component of the liposome.
 - 3. Liposomes according to either preceding claim, wherein the lipid component is one or more of phosphatidylcholine,
- cholesterol, phosphatidylglycerol, phosphatidylethanolamine, phosphatidylserine, phosphatidic acid, sphingomyelin or derivatives of these lipids.
 - 4. Liposomes according to claim 3, wherein the lipid components are phosphatidylcholine and cholesterol.
- 5. Liposomes according to any preceding claim, wherein the one or more substances are selected from carbohydrates and peptides.
- 6. Liposomes comprising, co-entrapped or covalently linked in the same liposome, a synthetic peptide and an adjuvant which potentiates the immunogenicity of the peptide.
 - 7. Liposomes according to claim 6, wherein the adjuvant is recognised by or interacts with T cells.
 - 8. Liposomes according to claim 6, wherein the adjuvant is a peptide having a T cell recognition site.
 - 9. Liposomes according to claim 6, wherein the adjuvant is IL-2.
- 10. Liposomes according to claim 6, wherein the adjuvant comprises a peptide having a T cell recognition site and 30 IL-2.
 - 11. Liposomes according to any of claims 5 to 10, wherein the peptide includes all or part of amino-acid residues 110-137 of Hepatitis B surface antigen.
- 12. Liposomes according to any preceding claim, for therapeutic use.

- 13. A vaccine composition comprising liposomes according to any preceding claim and a physiologically-acceptable excipient.
- 14. Use of liposomes according to any of claims 5 to 11, for the manufacture of a medicament for use in eliciting an immunogenic response to the synthetic peptide.
 - 15. A method for preparing liposomes having a mean diameter not exceeding 200 nm and containing one or more entrapped substances in an amount exceeding 10% of starting amounts.
 - 16. A method for producing liposomes having a mean diameter not exceeding 200 nm, comprising:
- (a) subjecting a mixture of lipid components and substances to be entrapped or covalently linked to the
 lipid component, to dehydration and controlled rehydration, to give large multilamellar vesicles; and
 - (b) subjecting the large multilamellar vesicles to microfluidisation.
- 17. A method according to claim 15 or claim 16, wherein 20 the starting concentration of lipid is no more than 100 μ mols per ml.
 - 18. Liposomes according to any of claims 15 to 17, wherein the total starting concentration of all substances is no more than 10 mg per ml.
- 25 19. A method according to any of claims 15 to 18, for preparing liposomes having the characteristics of any of claims 1 to 11.

FIGURE 1




FIGURE 2




FIGURE 3




FIGURE 4




FIGURE 5





FIGURE 6



	International Application No.	PC1/GB 91/0154/
CLASSIFICATION OF SUBJECT MATTER (If several classific	ication symbols apply, indicate all) ⁶	
According to International Patent Classification (IPC) or to both Na	itienal Classification and LPC	
Int.C1. 5 A61K9/127		•
L FIELDS SEARCHED		
Misleum	Documentation Sourches	
Chauffication System	Classification Symbols	
		···
nt.Cl. 5 A61K	•	•
	rd other than Minimum Documentation unexits are Included in the Fields Searched ⁸	
		<u> </u>
E. DOCUMENTS CONSIDERED TO BE RELEVANT		
ategory * Citation of Document, 31 with indication, where a	appropriate, of the relevant passages 12	Relevant to Claim Na.13
EP.A,0 290 296 (THE LIPOSO	ME COMPANY) 9 November	1,3,4,12
1988		2,5-11,
		13.14
	•	
VACCINE	· 	2
vol. 5, July 1987, GUILDFO	IRD	
pages 145 - 151; G.GREGORIADIS ET AL.: 'lip	acamae ae	
immunological adjuvants: a		
studies '		
see page 148, col.2, secon cited in the application)	d paragraph (document	
FR,A,2 624 741 (INSTITUT P	ACTEUD) 22 June 1080	5-10,13,
PR,A,2 024 /41 (10311101 P	ASTEUR) 23 Dulle 1909	14
see page 5 - page 6; examp	les 1,6	
	-/	
	•	ŀ
	. The base decrement will that after the late	manifest filling date
* Special categories of cited documents : 19 "A" document defining the general state of the art which is not	"I" inter document published after the inte or priority date and not in conflict with cited to understand the principle or the	the application but
omeidated to be of perfember reference	investisa	
"E" surilor document but published on er after the interactional filling date	"I" document of particular relevance; the consideral novel or cannot be	e considered to
"L" decreases which may throw doubts on priority claim(s) or which is cited to establish the publication date of another	involve an inventive step "Y" document of particular relovance; the e	isinal levertice
citation or other special reason (as specified) "O" successor referring to an oral dischause, use, exhibition or	councy be considered to involve an inv focusing to combined with one of more	milye step when the
Other meets	ments, such combination being obvious in the suc.	s to a person skilled
"P" decument published prior to the international filing date but later than the priority date claimed	"å" document manner of the same patent	haily
. CERTIFICATION		
ste of the Actual Completion of the Interactional Search	Date of Mailing of this International S	outch Report
•		
14 JANUARY 1992	30	JAN 1992
teractional Searching Authority	Signature of Authorized Officer	

Ė

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)						
ategory "	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No.				
	WO,A,9 004 412 (REGENTS OF THE UNIVERSITY OF MINNESOTA) 3 May 1990 see page 5, line 16 - line 31 see page 11, line 26 - page 13, line 22 see page 31 - page 32; example 6	11				
	US,A,4 235 877 (FULLERTON) 25 November 1980 see column 3 - column 4; examples 3,5	11				
	WO,A,9 004 943 (E. UNGER ET AL) 17 May 1990 see page 48 - page 51; examples 8,9	15-19				
·	•••					
·						

ANNEX TO THE INTERNATIONAL SEARCH REPORT ON INTERNATIONAL PATENT APPLICATION NO. GB 9101547 SA 51307

This armon lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the European Patent Office EDP file on

The European Patent Office is in no way liable for those particulars which are merely given for the purpose of information. 14/01/92

Putent decument cited in search report	Publication date	Potent family member(s)		Publication date	
EP-A-0290296	09-11-88	AU-A- AU-A- AU-A- JP-T- WO-A-	1481688 8141791 8141891 2502458 8806442	26-09-88 07-11-91 10-10-91 09-08-90 07-09-88	
FR-A-2624741	23-06-89	EP-A- WO-A-	0391962 8905631	17-10-90 29-06-89	
WO-A-9004412	03-05-90	AU-A- EP-A- CA-A-	4525589 0440742 2001737	14-05-90 14-08-91 19-01-91	
US-A-4235877	25-11-80	None			
WO-A-9004943	17-05-90	EP-A-	0442962	28-08-91	